## Practice 10

**PROBLEM** 1 Arithmetic

1. The prime factorization of 18<sup>2</sup> is \_\_\_\_\_\_.

2. Re-write  $p^r = w$  without a exponent function: \_\_\_\_\_

3. Simplify  $\frac{\log_3(7)}{\log_3(5)}$ : \_\_\_\_\_\_.

4. Re-write  $\log_2(16x^3)$  with no constants or operators in a log's argument:

5. What is  $\log_3(5) \log_5(3)$ ? \_\_\_\_\_.

**PROBLEM 2** Proof by Contradiction

Complete the following proof that  $\forall x \in \mathbb{Z}^+$ .  $(\log_3(x) \in \mathbb{Q}^+) \to (\exists n \in \mathbb{N} : x = 3^n).$ 

*Proof.* Assume that the implication does not hold; that is, that  $(\log_3(x) \in \mathbb{Q}^+) \land (\nexists n \in \mathbb{N} \cdot x = 3^n)$ . Since  $\log_3(x) \in \mathbb{Q}^+$ , there are positive integers a and b such that  $\log_3(x) = \frac{a}{b}$ . Re-writing that equation,

Since *a* and *b* are positive integers, both sides of the last equation above are integers. By the fundamental theorem of arithmetic, both sides must have the same prime factors, meaning that all of *x*'s factors must be 3. But that contradicts our assumption that  $\nexists n \in \mathbb{N}$ .  $x = 3^n$ .

Because the assumption led to a contradiction, it must be false; thus,

$$\left(\log_3(x) \in \mathbb{Q}^+\right) \to \left(\exists n \in \mathbb{N} : x = 3^n\right)$$

Want additional practice? Try the following: Simplify (show your work)

- $\log_3(5) + \log_3(2)$
- $\log_3(5) + \log_9(0.2)$
- $\log_3\left(\frac{5}{27}\right)$

$$\frac{\log_3(5)}{\log_2(7)}$$

• 
$$7\overline{\log_3(7)}$$

Complete

- $\log_{\sqrt{3}}(5) = \log_3\left( \begin{array}{c} \end{array} \right)$
- $\log_a(b)\log_a\left( \begin{array}{c} \end{array} \right) = 1$
- $\bullet \ \log_a(b) \log_b \left( \qquad \qquad \right) = 1$
- $\log_3(13) = \log_3(5) + \log_3\left($
- $3^{\log_5(7)} = 7^{\log_\Box(\Box)}$

Prove that

- there is no largest prime number. Use contradiction, with 1 + the product of all primes as part of how you get the contradiction.
- $(\log_a(b) = \log_b(a)) \rightarrow (a = b)$ . Both direct proof and contradiction should be able to work here.
- " $\forall n \in \{i \mid i \in \mathbb{Z} \land 1 < i < x\}$ .  $\log_i(x) \notin \mathbb{Q}$ " is true for all prime numbers x. Use contradiction.
- $3\log_2(10) < 10$ . Direct proof should be enough.
- $\log_3(10) > 2$ . Direct proof should be enough.